



## Step Therapy Criteria 2019

For information on obtaining an updated coverage determination or an exception to a coverage determination please contact Optimum HealthCare Member Services at 1-866-245-5360 or, for TTY/TDD users 711. From October 1 to March 31 from 8:00 am to 8:00 pm 7 days a week and April 1 to September 30 from 8:00 am to 8:00 pm Monday through Friday or visit [www.youroptimumhealthcare.com](http://www.youroptimumhealthcare.com).

For an indexed list of drugs please go to page 13.

# anti-emetics hrm

---

## Products Affected

- PHENADOZ SUPPOSITORY 12.5 MG RECTAL
- *promethazine hcl suppository 12.5 mg rectal*
- *promethazine hcl suppository 25 mg rectal*
- *promethazine hcl suppository 50 mg rectal*
- *promethazine hcl syrup 6.25 mg/5ml oral*
- *promethazine hcl tablet 12.5 mg oral*
- *promethazine hcl tablet 25 mg oral*
- *promethazine hcl tablet 50 mg oral*
- PROMETHEGAN SUPPOSITORY 25 MG RECTAL
- PROMETHEGAN SUPPOSITORY 50 MG RECTAL
- *trimethobenzamide hcl capsule 300 mg oral*

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY ONE OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Ondansetron, Granisetron. Step 2: Promethazine, Phenadoz, Promethagan, Trimethobenzamide. Criteria only applies for a diagnosis of nausea and vomiting. Criteria does not apply to the diagnosis of motion sickness. Step therapy only applies to members 65 years of age or older
-----------------	---

---

# anxiolytics hrm

---

## Products Affected

- *meprobamate tablet 200 mg oral*
- *meprobamate tablet 400 mg oral*

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY ONE OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Buspirone. Step 2: Meprobamate Step therapy only applies to members 65 years of age or older
-----------------	--

---

# atypicals

---

## Products Affected

- FANAPT TABLET 1 MG ORAL
- FANAPT TABLET 10 MG ORAL
- FANAPT TABLET 12 MG ORAL
- FANAPT TABLET 2 MG ORAL
- FANAPT TABLET 4 MG ORAL
- FANAPT TABLET 6 MG ORAL
- FANAPT TABLET 8 MG ORAL
- FANAPT TITRATION PACK TABLET 1 & 2 & 4 & 6 MG ORAL
- INVEGA SUSTENNA SUSPENSION 117 MG/0.75ML INTRAMUSCULAR
- INVEGA SUSTENNA SUSPENSION 156 MG/ML INTRAMUSCULAR
- INVEGA SUSTENNA SUSPENSION 234 MG/1.5ML INTRAMUSCULAR
- INVEGA SUSTENNA SUSPENSION 39 MG/0.25ML INTRAMUSCULAR
- INVEGA SUSTENNA SUSPENSION 78 MG/0.5ML INTRAMUSCULAR
- INVEGA TRINZA SUSPENSION 273 MG/0.875ML INTRAMUSCULAR
- INVEGA TRINZA SUSPENSION 410 MG/1.315ML INTRAMUSCULAR
- INVEGA TRINZA SUSPENSION 546 MG/1.75ML INTRAMUSCULAR
- INVEGA TRINZA SUSPENSION 819 MG/2.625ML INTRAMUSCULAR
- LATUDA TABLET 120 MG ORAL
- LATUDA TABLET 20 MG ORAL
- LATUDA TABLET 40 MG ORAL
- LATUDA TABLET 60 MG ORAL
- LATUDA TABLET 80 MG ORAL
- REXULTI TABLET 0.25 MG ORAL
- REXULTI TABLET 0.5 MG ORAL
- REXULTI TABLET 1 MG ORAL
- REXULTI TABLET 2 MG ORAL
- REXULTI TABLET 3 MG ORAL
- REXULTI TABLET 4 MG ORAL
- RISPERDAL CONSTA SUSPENSION RECONSTITUTED 12.5 MG INTRAMUSCULAR
- RISPERDAL CONSTA SUSPENSION RECONSTITUTED 25 MG INTRAMUSCULAR
- RISPERDAL CONSTA SUSPENSION RECONSTITUTED 37.5 MG INTRAMUSCULAR
- RISPERDAL CONSTA SUSPENSION RECONSTITUTED 50 MG INTRAMUSCULAR
- SAPHRIS TABLET SUBLINGUAL 10 MG SUBLINGUAL
- SAPHRIS TABLET SUBLINGUAL 2.5 MG SUBLINGUAL
- SAPHRIS TABLET SUBLINGUAL 5 MG SUBLINGUAL

## Details

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: aripiprazole, clozapine, risperidone, olanzapine, paliperidone, quetiapine, ziprasidone. Step 2: Fanapt, Invega Sustenna, Invega Trinza , Latuda, Rexulti, Risperdal Consta, Saphris. Step Therapy only applies to new starts only. Enrollees stabilized on medication will not be required to go through step therapy. Step therapy criteria only applies for the diagnosis of schizophrenia.
-----------------	---

# diabetes

---

## Products Affected

- AVANDIA TABLET 2 MG ORAL
- AVANDIA TABLET 4 MG ORAL

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Acarbose, Basaglar, Glimepiride, Glipizide, Glipizide ER, Glipizide XL, Glipizide/Metformin Hcl, Humalog, Humalog Mix 50/50, Humalog Mix 75/25, Humulin 70/30, Humulin N, Humulin R, Lantus, Lantus Solostar, Levemir, Levemir Flexpen, Metformin Hcl, Metformin Hcl Er, Nateglinide, Novolog, Novolog Flexpen, Novolog Mix 70/30, Novolin N, Novolin R, Novolin 70/30, Pioglitazone/Metformin, Repaglinide, Toujeo, Tresiba Step 2: Avandia.
-----------------	--

# dpp-4 inhibitors

---

## Products Affected

- JANUMET TABLET 50-1000 MG ORAL
- JANUMET TABLET 50-500 MG ORAL
- JANUMET XR TABLET EXTENDED RELEASE 24 HOUR 100-1000 MG ORAL
- JANUMET XR TABLET EXTENDED RELEASE 24 HOUR 50-1000 MG ORAL
- JANUMET XR TABLET EXTENDED RELEASE 24 HOUR 50-500 MG ORAL
- JANUMET XR TABLET EXTENDED RELEASE 24 HOUR 50-500 MG ORAL
- JANUVIA TABLET 100 MG ORAL
- JANUVIA TABLET 25 MG ORAL
- JANUVIA TABLET 50 MG ORAL
- KOMBIGLYZE XR TABLET EXTENDED RELEASE 24 HOUR 2.5-1000 MG ORAL
- KOMBIGLYZE XR TABLET EXTENDED RELEASE 24 HOUR 5-1000 MG ORAL
- KOMBIGLYZE XR TABLET EXTENDED RELEASE 24 HOUR 5-500 MG ORAL
- ONGLYZA TABLET 2.5 MG ORAL
- ONGLYZA TABLET 5 MG ORAL
- TRADJENTA TABLET 5 MG ORAL

## Details

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Glipizide/Metformin HCL, Metformin HCL, Metformin HCL ER, Pioglitazone/Metformin Step 2: Janumet, Januvia, Onglyza, Janumet XR, Kombiglyze, Tradjenta
-----------------	--

# glp1 agonist

---

## Products Affected

- BYETTA 10 MCG PEN SOLUTION PEN-INJECTOR 10 MCG/0.04ML SUBCUTANEOUS
- BYETTA 5 MCG PEN SOLUTION PEN-INJECTOR 5 MCG/0.02ML SUBCUTANEOUS
- TANZEUM PEN-INJECTOR 30 MG SUBCUTANEOUS
- TANZEUM PEN-INJECTOR 50 MG SUBCUTANEOUS
- VICTOZA SOLUTION PEN-INJECTOR 18 MG/3ML SUBCUTANEOUS

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Acarbose, Glimepiride, Glipizide, Glipizide ER, Glipizide XL, Glipizide/Metformin HCL, Metformin HCL, Metformin HCL ER, Pioglitazone/Metformin, Nateglinide, Repaglinide. Step 2: Tanzeum, Victoza, Byetta
-----------------	---

---

## non-sedating antihistamines

---

### Products Affected

- CLARINEX-D 12 HOUR TABLET  
EXTENDED RELEASE 12 HOUR 2.5-120  
MG ORAL
- *desloratadine tablet 5 mg oral*
- *desloratadine tablet dispersible 2.5 mg oral*
- *desloratadine tablet dispersible 5 mg oral*

### Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Allegra OTC, Allegra D OTC, Loratadine OTC, Loratadine D OTC, Cetirizine OTC, Cetirizine D OTC, Levocetirizine 0.5mg/ml oral soln., Levocetirizine 5mg OTC: Step 2: Desloratadine, Clarinex D. Only one drug, cetirizine, is required to be tried for the diagnosis of perennial allergic rhinitis.
-----------------	--

---



# ppi

---

## Products Affected

- *lansoprazole tablet dispersible 15 mg oral*
- *lansoprazole tablet dispersible 30 mg oral*

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: lansoprazole, lansoprazole OTC, Priolosec Otc, Nexium OTC, Omeprazole Otc, Omeprazole, Pantoprazole, Rabeprazole, Prevacid OTC, Zegerid OTC. Step 2: lansoprazole Solutabs. For the diagnosis of risk reduction of NSAID associated gastric ulcer only lansoprazole is required.
-----------------	---

---

## renin inhibitors

---

### Products Affected

- TEKTURNA HCT TABLET 150-12.5 MG ORAL
- TEKTURNA HCT TABLET 150-25 MG ORAL
- TEKTURNA HCT TABLET 300-12.5 MG ORAL
- TEKTURNA HCT TABLET 300-25 MG ORAL
- TEKTURNA TABLET 150 MG ORAL
- TEKTURNA TABLET 300 MG ORAL

### Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS, 1 DRUG FROM EACH CLASS, IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: ACE-Inhibitors (including combinations with HCTZ) - Benazepril Hcl, Benazepril Hctz, Captopril, Captopril /Hctz, Enalapril Maleate, Enalapril Maleate/Hctz, Fosinopril Sodium, Fosinopril sodium/Hctz, Lisinopril, Lisinopril /Hctz, Quinapril Hcl, Quinipril/HCTZ, Trandolapril, Ramipril. ARBs (including combinations with HCTZ) - Olmasartan, Olmasartan/Hct, losartan, losartan/HCT, irbesartan, irbesartan/Hctz, valsartan, valsartan/Hctz. Step 2: Tekturna, Tekturna Hct
-----------------	--

---

# symlin

---

## Products Affected

- SYMLINPEN 120 SOLUTION PEN-INJECTOR 2700 MCG/2.7ML  
SUBCUTANEOUS
- SYMLINPEN 60 SOLUTION PEN-INJECTOR 1500 MCG/1.5ML  
SUBCUTANEOUS

## Details

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Basaglar, Humalog, Humalog Mix 50/50, Humalog Mix 75/25, Humulin 70/30, Humulin N, Humulin R, Lantus, Levemir, Novolog, Novolog Mix 70/30, Novolin R, Novolin N, Novolin 70/30, Toujeo, Tresiba Step 2: Symlin
-----------------	---

# topical\_immunomodulators

---

## Products Affected

- ELIDEL CREAM 1 % EXTERNAL
- *tacrolimus ointment 0.03 % external*
- *tacrolimus ointment 0.1 % external*

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF ANY TWO OF THE FOLLOWING DRUGS IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Ala-Cort, Alclometasone Dipropionate, Amcinonide, Augmented Betamethasone Dipropionate, Betamethasone, Betamethasone Dipropionate, Betamethasone Valerate, Clobetasol Propionate, Clobetasol Propionate Emollient, Desonide, Desoximetasone, Diflorasone Diacetate, Fluocinolone Acetonide, Fluocinonide, Fluticasone, Halobetasol Propionate, Hydrocortisone Butyrate, Hydrocortisone Valerate, Mometasone Furoate, Prednicarbate, Triamcinolone Acetonide, Triamcinolone Acetonide In Absorbase. Step 2: Elidel, Tacrolimus
-----------------	--

# ULORIC

---

## Products Affected

- ULORIC TABLET 40 MG ORAL
- ULORIC TABLET 80 MG ORAL

## Details

---

<b>Criteria</b>	Step 1- PATIENT NEEDS TO HAVE A DOCUMENTED TRIAL OF THE FOLLOWING DRUG/S IN THE PREVIOUS 180 DAYS BEFORE MOVING TO STEP 2: Allopurinol. STEP 2: Uloric
-----------------	--

---

**Index (Índice)**

AVANDIA TABLET 2 MG ORAL.....	4	JANUMET XR TABLET EXTENDED	
AVANDIA TABLET 4 MG ORAL.....	4	RELEASE 24 HOUR 100-1000 MG	
BYETTA 10 MCG PEN SOLUTION		ORAL.....	5
PEN-INJECTOR 10 MCG/0.04ML		JANUMET XR TABLET EXTENDED	
SUBCUTANEOUS.....	6	RELEASE 24 HOUR 50-1000 MG ORAL....	5
BYETTA 5 MCG PEN SOLUTION PEN-		JANUMET XR TABLET EXTENDED	
INJECTOR 5 MCG/0.02ML		RELEASE 24 HOUR 50-500 MG ORAL.....	5
SUBCUTANEOUS.....	6	JANUVIA TABLET 100 MG ORAL.....	5
CLARINEX-D 12 HOUR TABLET		JANUVIA TABLET 25 MG ORAL.....	5
EXTENDED RELEASE 12 HOUR 2.5-		JANUVIA TABLET 50 MG ORAL.....	5
120 MG ORAL.....	7	KOMBIGLYZE XR TABLET	
<i>desloratadine tablet 5 mg oral.....</i>	<i>7</i>	EXTENDED RELEASE 24 HOUR 2.5-	
<i>desloratadine tablet dispersible 2.5 mg</i>		1000 MG ORAL.....	5
<i>oral.....</i>	<i>7</i>	KOMBIGLYZE XR TABLET	
<i>desloratadine tablet dispersible 5 mg oral.....</i>	<i>7</i>	EXTENDED RELEASE 24 HOUR 5-1000	
ELIDEL CREAM 1 % EXTERNAL.....	11	MG ORAL.....	5
FANAPT TABLET 1 MG ORAL.....	3	KOMBIGLYZE XR TABLET	
FANAPT TABLET 10 MG ORAL.....	3	EXTENDED RELEASE 24 HOUR 5-500	
FANAPT TABLET 12 MG ORAL.....	3	MG ORAL.....	5
FANAPT TABLET 2 MG ORAL.....	3	<i>lansoprazole tablet dispersible 15 mg oral....</i>	<i>8</i>
FANAPT TABLET 4 MG ORAL.....	3	<i>lansoprazole tablet dispersible 30 mg oral....</i>	<i>8</i>
FANAPT TABLET 6 MG ORAL.....	3	LATUDA TABLET 120 MG ORAL.....	3
FANAPT TABLET 8 MG ORAL.....	3	LATUDA TABLET 20 MG ORAL.....	3
FANAPT TITRATION PACK TABLET 1		LATUDA TABLET 40 MG ORAL.....	3
& 2 & 4 & 6 MG ORAL.....	3	LATUDA TABLET 60 MG ORAL.....	3
INVEGA SUSTENNA SUSPENSION 117		LATUDA TABLET 80 MG ORAL.....	3
MG/0.75ML INTRAMUSCULAR.....	3	<i>meprobamate tablet 200 mg oral.....</i>	<i>2</i>
INVEGA SUSTENNA SUSPENSION 156		<i>meprobamate tablet 400 mg oral.....</i>	<i>2</i>
MG/ML INTRAMUSCULAR.....	3	ONGLYZA TABLET 2.5 MG ORAL.....	5
INVEGA SUSTENNA SUSPENSION 234		ONGLYZA TABLET 5 MG ORAL.....	5
MG/1.5ML INTRAMUSCULAR.....	3	PHENADOZ SUPPOSITORY 12.5 MG	
INVEGA SUSTENNA SUSPENSION 39		RECTAL.....	1
MG/0.25ML INTRAMUSCULAR.....	3	<i>promethazine hcl suppository 12.5 mg</i>	
INVEGA SUSTENNA SUSPENSION 78		<i>rectal.....</i>	<i>1</i>
MG/0.5ML INTRAMUSCULAR.....	3	<i>promethazine hcl suppository 25 mg rectal....</i>	<i>1</i>
INVEGA TRINZA SUSPENSION 273		<i>promethazine hcl suppository 50 mg rectal....</i>	<i>1</i>
MG/0.875ML INTRAMUSCULAR.....	3	<i>promethazine hcl syrup 6.25 mg/5ml oral.....</i>	<i>1</i>
INVEGA TRINZA SUSPENSION 410		<i>promethazine hcl tablet 12.5 mg oral.....</i>	<i>1</i>
MG/1.315ML INTRAMUSCULAR.....	3	<i>promethazine hcl tablet 25 mg oral.....</i>	<i>1</i>
INVEGA TRINZA SUSPENSION 546		<i>promethazine hcl tablet 50 mg oral.....</i>	<i>1</i>
MG/1.75ML INTRAMUSCULAR.....	3	PROMETHEGAN SUPPOSITORY 25	
INVEGA TRINZA SUSPENSION 819		MG RECTAL.....	1
MG/2.625ML INTRAMUSCULAR.....	3	PROMETHEGAN SUPPOSITORY 50	
JANUMET TABLET 50-1000 MG ORAL....	5	MG RECTAL.....	1
JANUMET TABLET 50-500 MG ORAL....	5	REXULTI TABLET 0.25 MG ORAL.....	3
		REXULTI TABLET 0.5 MG ORAL.....	3

REXULTI TABLET 1 MG ORAL.....	3	<i>trimethobenzamide hcl capsule 300 mg</i>	
REXULTI TABLET 2 MG ORAL.....	3	<i>oral</i> .....	1
REXULTI TABLET 3 MG ORAL.....	3	ULORIC TABLET 40 MG ORAL.....	12
REXULTI TABLET 4 MG ORAL.....	3	ULORIC TABLET 80 MG ORAL.....	12
RISPERDAL CONSTA SUSPENSION RECONSTITUTED 12.5 MG INTRAMUSCULAR.....	3	VICTOZA SOLUTION PEN-INJECTOR 18 MG/3ML SUBCUTANEOUS.....	6
RISPERDAL CONSTA SUSPENSION RECONSTITUTED 25 MG INTRAMUSCULAR.....	3		
RISPERDAL CONSTA SUSPENSION RECONSTITUTED 37.5 MG INTRAMUSCULAR.....	3		
RISPERDAL CONSTA SUSPENSION RECONSTITUTED 50 MG INTRAMUSCULAR.....	3		
SAPHRIS TABLET SUBLINGUAL 10 MG SUBLINGUAL.....	3		
SAPHRIS TABLET SUBLINGUAL 2.5 MG SUBLINGUAL.....	3		
SAPHRIS TABLET SUBLINGUAL 5 MG SUBLINGUAL.....	3		
SYMLINPEN 120 SOLUTION PEN- INJECTOR 2700 MCG/2.7ML SUBCUTANEOUS.....	10		
SYMLINPEN 60 SOLUTION PEN- INJECTOR 1500 MCG/1.5ML SUBCUTANEOUS.....	10		
<i>tacrolimus ointment 0.03 % external</i> .....	11		
<i>tacrolimus ointment 0.1 % external</i> .....	11		
TANZEUM PEN-INJECTOR 30 MG SUBCUTANEOUS.....	6		
TANZEUM PEN-INJECTOR 50 MG SUBCUTANEOUS.....	6		
TEKTURN HCT TABLET 150-12.5 MG ORAL.....	9		
TEKTURN HCT TABLET 150-25 MG ORAL.....	9		
TEKTURN HCT TABLET 300-12.5 MG ORAL.....	9		
TEKTURN HCT TABLET 300-25 MG ORAL.....	9		
TEKTURN TABLET 150 MG ORAL.....	9		
TEKTURN TABLET 300 MG ORAL.....	9		
TRADJENTA TABLET 5 MG ORAL.....	5		